

**ErbB 2****PLD-608****Package :** 可询价**Product Name :** ErbB 2**Cat.No.:** PLD-608**Synonyms :** NEU; NGL; HER2; TKR1; CD340; HER-2; MLN 19; HER-2/neu**Application :** IHC-P**Reactivity :** Human**Host species :** Rabbit**Background**

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq, Jul 2008]

**Immunogen**

Recombinant protein of human ErbB 2

**Gene ID**

2064

**Swiss Prot**

P04626

**Synonyms**

NEU; NGL; HER2; TKR1; CD340; HER-2; MLN 19; HER-2/neu

**Reactivity**

Human

**Application**

IHC-P

**Recommended dilution**

IHC-P: 1:5000

**Host species**

Rabbit

**Clonality**

Monoclonal

**Isotype**

IgG

**Purity**

Affinity Purification

**Conjugation**

Un-conjugated

**Storage Stability**

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.